

A Map from Scalar Field Theory to Integer Polynomial Solutions

Gordon Chalmers

e-mail: gordon@quartz.shango.com

Abstract

The terms in the quantum scattering in scalar field theory models is parameterized by the invariants $\prod s_{ij}^{n_{ij}}$. The s_{ij} are kinematic two-particle invariants, and the n_{ij} are integers. The coefficients of these terms are computed via integrating all Feynman diagrams, or within the derivative expansion by solving the iteration equations. The latter has been provided recently; the functions which are prefactors of the individual terms $\prod s_{ij}^{n_{ij}}$ can be interpreted as terms in the expansions of L-series, which may be specified by collections of their zeroes. Once finding the appropriate elliptic curve coefficients, these quantum field solutions provide an algorithm to determining all of the mod p zeros to the algebraic curves. The latter is presumably determined by 'experimental' computer modeling or by the appropriate determination of the quantum prefactors.

The quantum solution to scalar field models is determined by the initial conditions of the bare Lagrangian. Given the initial conditions, there is an algorithm for determining the quantum scattering. These initial conditions are examined, together with solving all of the integrals, for ϕ^3 theory in [1]. The solutions to the terms in the effective action, or scattering, is developed in prior works.

The bare Lagrangian is contained in the expansion,

$$\mathcal{L} = \frac{1}{2}\phi\Box\phi + \frac{1}{2}m^2\phi^2 + \frac{1}{3!}\lambda_3\phi^3 \quad (1)$$

with the higher order terms

$$\mathcal{L}_{\text{bare}} = \frac{1}{4!}\phi^4 + \sum \mathcal{O}_i . \quad (2)$$

The operators $\mathcal{O}_i$ may contain terms such as $\Box\phi^2\Box\phi^2$ and all combinations of derivatives with the fields ϕ . The initial theory in (1) and (2).

The solution to these scalar models requires the summation of terms, i.e. rainbow graphs, illustrated in [1]. These graphs were integrated in [1], and the integration leads to a simplification of the arbitrary loop graphs in the conventional Feynman diagrams. The summation appears complicated due to the combinatorics of the tree diagrams obtained from the theory in (1) and (2); however, there are simplifications that allow the summation to be performed [2] in the arbitrary bare Lagrangian described.

The scattering at n -point in these *scalar* models is described by functions in k -space,

$$A_{s_{ij}} = f_{s_{ij}, n_{ij}} \prod s_{ij}^{n_{ij}} , \quad (3)$$

with the invariants $s_{ij} = (k_i + k_j)$ and with the n_{ij} integers labeling the expansion. The set of all combinations spans the complete scattering at an arbitrary n -point (with n labeling the k_i in the product). The functions f are functions of the couplings λ_3, λ_4 , as well as the coefficients of the higher derivative terms in the bare theory as spanned by the operators $\mathcal{O}_i$.

The determination of the functions $f_{s_{ij}, n_{ij}}$ are given in [1]. It is remarkable that a closed form solution can almost be given, in [1], for an arbitrary bare Lagrangian. (As mentioned, the summations in [1] can also be obtained, most likely, in [2].)

The determination of the functions $f_{s_{ij}, n_{ij}}$ could require a different determination, for example requiring the period matrix as in the case of the $N = 2$ superpotential (or prepotential). However, in the scalar field models there are an infinite number of functions spanned by the non-two derivative terms.

These functions are a power series in the coupling λ_3 , and in the other couplings. If all of the couplings are chosen to be proportional to λ_3 , with factors of the mass m (or in a conventional cutoff setting involving Λ), then these functions are arbitrary. This would involve power series possibly multiplying the bare terms. However, the theory involving only the λ_3 term is sufficient for this analysis.

The series of functions $f_{s_{ij}, n_{ij}}$ span a set of determined functions, all of which in perturbation theory are power series in the coupling λ (i.e. in ϕ^3 , ϕ^4 , or with the improvements).

These functions can be turned into a collection of zeros. For example, these functions, as

$$f_i = \sum a_{i,j} \lambda^j \quad (4)$$

can be converted into the zero set,

$$f_i = \prod (\tilde{\lambda}_{i,a} + \lambda_a) \quad (5)$$

with the factors $\tilde{\lambda}_{i,a}$ complex variables. The i labels the various terms in the quantum scattering at n -point. These functions can be interpreted as the zero set to an associated L-series

$$L(C, \lambda) = \prod (1 - a_p p^{-2\lambda} + p^{1-2\lambda})^{-1} . \quad (6)$$

These L-series have an arbitrary set of zeros on the complex plane, given the parameters $a_p = N_p + p$ which might be arbitrary; however, the curve and its solutions

$$y^2 = x^3 + ax + b \bmod p \quad (7)$$

with p prime generate these solutions; N_p counts the solutions at a given prime p . The numbers a and b which determine these solutions could be large or small. These

L-series have an expansion, in λ which is a power series; they also are described by the zero set of the function as in (5) (the simplest example is the Riemann zeta function).

The interpretation of the quantum scattering, as described in the scalar field theory [1] and in [2], in terms of the elliptic function is valuable for a number of reasons. Primarily, once the curves' numbers a_i and b_i are determined, then all of the solutions' countings a_p in (6) can be found by an inverse transform; the solution to the a_i and b_i curve parameters depends on the n -point number and the set of integers n_{ij} (defining $\prod s_{ij}^{n_{ij}}$). Given the parameters a_i and b_i , this solution can be obtained, of which there no known recipe is known.

The terms in the effective action has been explained for an arbitrary tree solution in [1] and [2]. Apart from a solution based on the integrated terms as generated in a general setting [1], the symmetries of these tree diagrams is important to find the parameters a_i and b_i for the general term $\prod s_{ij}^{n_{ij}}$.

The parameters a_i and b_i require a different determination of the functions f_i from the explicit integral solution based on the derivative expansion. Once these terms are described, for a potential arbitrary set of conditions in the bare theory, which would generate an arbitrary curve (with a_i and b_i), then any elliptic curve could be modeled. This elliptic curve modeling would generate all of the solution numbers a_p in (6).

This work has bearing on the base interpretation of the elliptic modeling in [3], the gauge and gravity quantum solution (with [4]), and the compact expressions for gauge amplitudes in [5].

References

- [1] G. Chalmers, *Quantum Solutions to Scalar Field Theory Models*, physics/050518.
- [2] G. Chalmers, *in progress*.
- [3] G. Chalmers, *Integer and Rational Solutions to Polynomial Equations*, physics/0503200.
- [4] G. Chalmers, *Quantum Gauge Theory Amplitude Solutions*, physics/0505077.
- [5] G. Chalmers, *Very Compact Expressions for Amplitudes*, physics/0502058.